The Journal of Bone and Joint Surgery

ANNOTATION

SURGERY OF THE HAND

In 1916 Biesalski and Mayer* described their experience of tendon transplantation in animals and in the same year Mayer† published a paper on the physiological method of tendon transfer. Two years later Bunnell‡ described the operation of tendon repair in the finger. From these beginnings and from the inspiration of such men as Bunnell, Kanavel, Koch and Mayer has arisen the wide and increasing interest shown in surgery of the hand. Bunnell's monumental work, in particular, has laid the solid foundation upon which the younger generation is building. In most countries of the world there are now small groups of surgeons developing this long neglected surgical field and achieving a high standard of result. The hand has become the focal point of interest in which the two specialities of orthopaedic and plastic surgery meet. A new body of surgeons is arising who, while maintaining their interest in general orthopaedic or plastic work, have a special interest in the hand. Great Britain is sharing in this advance, but much more requires to be accomplished if this country is to remain in the front line of progress.

Reconstruction of the hand, "the jewelry of orthopaedic surgery," as Bunnell called it, does not favour the system of combined operation that is accepted and indeed necessary in general limb surgery. It is unwise for more than one surgeon to work upon the complex anatomy of the hand, and thus create a division of responsibility, and those who practise hand surgery must be capable of all the duties of this work. There is no place here for the surgeon who can repair the skin but not the nerves, or the surgeon who can suture tendons but not restore skin defects. This new speciality not only demands a particular type of training and a high degree of craftsmanship, but it is probably true to say that it calls for a particular temperament, perhaps somewhat like that of a small boy who takes his toy to pieces in order to see how it works, and then has the patience to put it together again!

Several centres for hand surgery have been formed by individual effort in this country, and the patient with the crippled hand does not need to travel far to reach a service where his problem is not exceptional. Unhappily, it is a very different story for the patient who goes to the casualty department of many general hospitals. He may be cared for by a member of the junior staff who has had little, if any, special training. The operating theatres are too busy with major surgical work to allow for the intrusion of such a case, still classified as minor, but which may need as much care and skill as the abdominal emergencies which jeopardise life. The hand is unique in its combination of specialised tissues, the skin adapted for many forms of sensibility, the delicate gliding parts for precise movements and the powerful muscles and tendons for gripping. These tissues react badly to the oedema which follows injury, and

* BIESALSKI, K., and MAYER, L. (1916): Die physiologische Sehnenverpflanzung. Berlin: J. Springer.

† MAYER, L. (1916): The Physiological Method of Tendon Transplantation. I. Historical; Anatomy and Physiology of Tendons. II. Operative Technique. III. Experimental and Clinical Experiences. Surgery, Gynecology and Obstetrics, **22**, 182; 298; 472.

‡ BUNNELL, S. (1918): Repair of Tendons in the Fingers and Description of Two New Instruments. Surgery, Gynecology and Obstetrics, **26**, 103.

the joint stiffness which so quickly appears is most difficult to overcome once it is established. Wounds, apparently trivial, may divide a tendon or nerve and lead to months of incapacity. Wounds that are left open to granulate heal with deep fibrosis and skin contractions which could have been avoided by the use of an immediate free skin graft or a suitably fashioned flap.

Rank and Wakefield* have recently drawn attention to the magnitude of the problem. In 1949, 30 per cent of all the injury benefits paid under the National Insurance Act of Great Britain for accidents and prescribed diseases were for injuries of the hand. Every year in the United States of America there are approximately 2,000,000 disabling work injuries and some 75 per cent affect the hand. In the State of Victoria industrial injuries reach a yearly average of 5,350, and, of this total, 1,950 consist of hand injuries.

The hand is one of man's most precious possessions, and the perfect co-ordination between brain and hand has determined man's destiny, and his place in nature. This symbol of man's power, this instrument of his perception, this mirror of his emotion; what is more versatile than the human hand, and what tool more necessary for the support of life? Great is the disaster when these attributes are impaired or lost by injury and disease.

There is need of a more acute sense of our responsibility for the care of these very common injuries and for an enthusiasm like that which led to the establishment of our fracture clinics. There are two ways in which the challenge can be met. Firstly by the formation of hand units attached to every accident service, and secondly by the training of young surgeons in this work, whether their determined career be orthopaedic or plastic surgery.

R. G. PULVERTAFT.

* RANK, B. K., and WAKEFIELD, A. R. (1953): *Surgery of Repair as Applied to Hand Injuries*, 3. Edinburgh and London: E. & S. Livingstone Ltd.