The Journal of Bone and Joint Surgery

EDITORIALS AND ANNOTATIONS

TREATMENT OF TUBERCULOSIS OF THE SPINE

Evaluation of the methods of treatment of skeletal tuberculosis is both necessary and difficult at this time—necessary because we are now approaching the time when the value of antibiotics may be reasonably assessed, and difficult because of the many factors to be taken into account, such as the age of the patient, the site of the disease and even the part of the country.

There seems to be reliable evidence that the primary mortality in bone and joint tuberculosis has been reduced by the use of antibiotics. If this be accepted, then recurrence of disease assumes an even greater importance in the assessment of results than it had in the past. Changes in the accepted methods of treatment should await reliable evidence not only that the mortality rate is less, but that morbidity is reduced through less time in hospital and fewer recurrences.

Direct attack on a focus of tuberculous disease has been an accepted method of approach when it was—or appeared to be—possible to excise completely the diseased organ or tissue. When removal was not possible the indications for early surgery were less clearly defined and less generally accepted. The risks of general dissemination of the disease and of the formation of sinuses were strong deterrents to adventurous surgery. These two possible complications, which were perhaps over-emphasised in the past, may now be less important since the introduction of the antibiotics. Direct attack on spinal disease by the antero-lateral approach was first carried out by Menard and others towards the end of the last century. These procedures were principally concerned with drainage of the diseased site, and the results were such that the operation was abandoned. Ito *et al.* (1934) reported ten patients treated by partial or complete excision of thoracic vertebrae by the costotransversectomy approach, and of lumbar vertebrae by the extraperitoneal approach. All the patients survived the operations, but sufficient time had not elapsed at the time of writing to assess the results. Since then no further report on the condition of these patients has appeared. No reports of other series treated by this method have been discovered since then except that by Felländer quoted below. I understand that the method has been used by Kastert of Stuttgart and that some results have recently been published—though it has not been possible to obtain a copy of this publication.

The results of treatment by conservative measures, including spinal fusion after the disease has been arrested, leave much to be desired, but they have been better than those produced by any other method up to the present. The principal drawback has been the long period of treatment required, with its inevitable mental and physical trauma.

Anything that can be done to decrease the length of treatment and still produce equally good short-term and long-term results is to be welcomed. It is perhaps wise to recall that, owing to the chronic nature of the disease, it is present in the spine for a year or more before diagnosis is possible, and that the safety of the patient—so far as the future is concerned when treatment is completed—depends as much on the general as on the local resistance to the disease. Much of the generally accepted regime of conservative treatment has been based on these two points.

In this number of the Journal Wilkinson mentions some published results of conservative treatment. The two series published by Girdlestone to which he refers were designed to compare the results of conservative treatment alone with those of conservative treatment followed by grafting when the disease was arrested. In the latter group 65 per cent were well and working after five years. Of the 15 per cent not mentioned by Wilkinson 8 per cent had been excluded because it was not possible to trace them, and 7 per cent because their treatment had not been completed five years before the follow-up was undertaken. It was established that none of the 8 per cent had died. The figure of 65 per cent was therefore a conservative estimate, and might have been increased by some of the missing 15 per cent. This review was carried out twenty years ago.

In Wilkinson's series the number of cases in each group is too small to bear statistical analysis, and, although they are grouped together in the summary from the point of view of results, no composite analysis is given of the length of follow-up, length of stay in hospital, further operations, etc. For instance, of the thirty-four adults only four had been followed up for five years. If bony ankylosis is accepted as the criterion of a satisfactory result in established spinal tuberculosis, the proportion of those in which a satisfactory result was obtained was 55.4 per cent.

Felländer (1955) has recently produced a complicated analysis of 100 patients in which surgical attack on the vertebral lesion was carried out. This paper demonstrates very well the difficulty of subjecting all the various factors to analysis. For example, in thirty-nine patients spinal fusion had been carried out before the radical operation, and in twenty-six it was undertaken after the radical operation. These sixty-five patients are compared unfavourably with thirty-six in which no spinal fusion followed the radical procedure. The thirty-nine patients represented a group in which the conservative treatment had failed to arrest the disease. No comparison is made of the twenty-six grafted after the radical operation with the thirty-six in which no graft was carried out. In this group of 100 no less than thirty-six had recurrence of sinuses or abscesses after the radical procedure. In thirty of these patients forty additional operations were carried out in relation to the recurrence of disease. The average length of hospital treatment was 169 ± 12 days. The period of observation was one to four years. Sixty-two patients were fully able to work, and seventy-three could be regarded as clinically healed with respect to the spondylitis. The mortality rate in the group during the period of observation was 9 per cent, three deaths being directly related to the operation. No brief comment on this excellent analysis can do it justice, and those interested in this subject are advised to read it in full.

We are all searching for steps that will shorten the long period of treatment required, if it continues to be based on conservative measures designed to produce good general health, arrest and quiescence of the local lesion and a strong and stable spine at the site of the disease. It may be that the direct attack on the spinal lesion by removal of diseased tissue, and improvement in blood supply to the remaining bone, will provide a step in this direction. But the evidence that this method decreases morbidity and leads to better long-term results is as yet insufficient to justify its general acceptance.

J. C. SCOTT.

REFERENCES

- FELLÄNDER, M. (1955): Radical Operation in Tuberculosis of the Spine. *Acta Orthopaedica Scandinavica*, Supplementum No. 19.
 ITO, H., TSUCHIYA, J., and ASAMI, G. (1934): A New Radical Operation for Pott's Disease. Report of Ten Cases. *Journal of Bone and Joint Surgery*, **16**, 499.
 WILKINSON, M. C. (1953): Synovectomy and Curettage in the Treatment of Tuberculosis of Joints. *Journal of Bone and Joint Surgery*, **35-B**, 209.