RECHERCHES SUR LA CONSTITUTION DE L'OS ADULTE. By J. VINCENT, Boursier de l'Institut Interuniversitaire des Sciences Nucléaires. $9\frac{1}{2} \times 6\frac{1}{4}$ in. Pp. 153, with 41 figures. 1955. Bruxelles: Éditions Arscia. Price Belgian francs 200.

In recent years much has been added to the knowledge of bone structure by the application of new techniques. Histochemical methods have supplemented the classical histological studies, and microradiography and autoradiography in particular have shed light on the physiological processes concerned in bone formation and bone resorption. Microradiography consists of the determination

FIG. 2

Microradiograph ($\times 135$) of a transverse section of the radial diaphysis of an adult dog. The dark dots are the lacunae containing osteocytes and the varying shades of grey represent different degrees of calcification. *a* and *b* are resorption cavities. *c*, *d* and *e* are completely formed, partially calcified osteones.

FIG. 3

Comparison of this photomicrograph ($\times 135$) of the decalcified unstained section and its microradiograph above shows uncalcified borders of "preosseous tissue" around the resorption cavities.

on a microscopic scale of the x-ray absorption of thin slices of tissue. A thin section of undecalcified bone is placed in contact with a fine-grained photographic emulsion and exposed to soft x-rays. After processing, the radiographic image can be examined under the microscope, and, from the variations in x-ray absorption, inferences can be drawn about the degree of mineralisation in the different areas of the preparation. Engstrom and his colleagues in the Department of Physical Cell Research at the Karolinska Institute in Stockholm have published much work in this field. Autoradiography is the study of the tissue uptake of radioactive substances, as shown by their effect on a photographic emulsion placed in contact with the material to be examined. Work on bone

mineralisation using lead isotopes was carried out by Lomholt as early as 1930, while in recent years the more physiological elements, radiophosphorus P^{32} , and radiocalcium Ca^{45} , have been used for this purpose, and studies of the uptake of C^{14} bicarbonate, and S^{35} sulphate, have also thrown light on the formation of the organic matrix of bone.

The present monograph has the form of a postgraduate dissertation. It is from Professor Lacroix's laboratory at Louvain in Belgium, and correlates histological, microradiographic and autoradiographic studies on the structure of bone. The techniques that have been used are described fully enough to be of use to other workers. The author is chiefly concerned with the bone of the adult dog, and a short section on human adult bone serves chiefly to support the choice of the dog as the experimental animal and to point out a few minor structural differences. The main interest in this work is the light it throws on Haversian remodelling in cortical bone. Dr Vincent has been able to show the steps that occur in the formation of new osteones as they come to replace resorption cavities.

"Pre-osseous" tissue (the "osteoid" tissue of English writers) is formed first. It is uncalcified and so is not demonstrable in microradiographs but is seen in histological preparations. In alcohol-fixed material decalcified by "versene" it stains orthochromatically with toluidin blue. The initially formed tissue shows uptake of radioactive sulphur, but not of radioactive calcium. Later, this "pre-osseous" tissue becomes calcified, rapidly acquiring about two-thirds of its ultimate content of bone salt. It is then apparent in microradiographs, and shows active uptake of radioactive calcium. The calcified tissue stains metachromatically with toluidin blue. The final completion of the mineralisation of the osteone is a slow process.

By serial studies after the administration of radioactive sulphur Dr Vincent has been able to estimate that in the adult dog the deposition of an osteone takes about six weeks but its complete calcification takes more than twelve weeks. He has found also that Haversian remodelling is an intermittent process and may cease temporarily either locally or in the skeleton as a whole. In spongy bone interpretation is more difficult, but here also there is evidence of remodelling. This occurs particularly in the metaphysis and resembles that seen in cortical bone.

Finally Dr Vincent discusses his results in relation to the general background of bone physiology and biochemistry, and suggests further lines of inquiry which appear capable of solution by the methods of study outlined.

The work is effectively illustrated by sets of comparable photomicrographs, microradiographs and autoradiographs, which add much to the clarity of the exposition. It is unfortunate that these illustrations are grouped together at the end of the book and not placed at the points in the text where they are discussed. There is an excellent bibliography.—Mary CATTO.

MANAGEMENT OF LIFE-THREATENING POLIOMYELITIS. Copenhagen, 1952-1956. Edited by H. C. A. LASSEN, M.D., Professor of Epidemiology, University of Copenhagen; Chief Physician, Department of Communicable Diseases, Blegdamshospitalet, Copenhagen; with twenty-one other contributors. $8\frac{1}{2} \times 5\frac{1}{2}$ in. Pp. xi+179, with 55 figures and 32 tables. 1956. Edinburgh and London: E. & S. Livingstone Ltd. Price 22s. 6d.

This book reports an epidemic of poliomyelitis which was in many respects the worst on record. The incidence in Copenhagen reached 105 paralytic and 153 non-paralytic patients per 100,000 population. At the peak there were 320 hospital admissions in a week. Of a total of 5,675 patients, 345 had severe paralysis of respiratory muscles or of swallowing and of these 42 per cent died. Of the 345, twelve had failure of swallowing alone, 254 respiratory muscle failure and no less than 79 had respiratory and pharyngo-laryngeal paralysis. The Danes had to treat these desperately ill patients with no hope of providing negative pressure cabinet respirators for more than a negligible fraction; indeed, this group of patients included an unheard of proportion for whom that type of aid would have been ineffective. A team of physicians, anaesthetists and laryngologists therefore improvised a method and recruited assistants. This team aided by neurologists, epidemiologists, biochemists and pathologists has now given us its results.

A method providing adequate ventilation had to be devised with which patency of the air passages could also be maintained. Postural drainage and pharyngeal suction had been proved completely inadequate for patients with secretions in the tracheobronchial tree resulting from failure of swallowing or from anoxic and hypercapnic hypersecretion. The method had also to allow both adequate oxygenation and carbon dioxide removal. The need was met by tracheotomy and insertion of an inflatable cuffed anaesthetic tube. Positive pressure ventilation was achieved by the manual compression of a rubber bag attached to this tube; the circuit included both a soda-lime carbon dioxide absorber and a supply of equal parts of oxygen and nitrogen. Frequent aspiration of the trachea and main bronchi was possible through the tracheotomy. Teams of helpers worked in shifts with this apparatus