of all the factors will sometimes make amputation of the digit the best primary treatment because patients show a remarkable reluctance later to have a stiff, livid, painful and useless appendage removed. This is an attitude dictated not only for reasons of compensation but because of the natural revulsion against mutilation, as the writer has emphasised in his opening sentence.

Let us then remember that increasingly reconstructive surgery includes, as much or more than amputation, the restoration of limbs which though not always completely normal have that boon of mankind—the abilities to feel and to grasp with our hands and to reach and to feel the ground with our feet.

H. OSMOND-CLARKE.

REFERENCE

GILLIS, L. (1956): Amputations in Children. *Annals of the Royal College of Surgeons of England*, **19**, 335.

OSTEOCHONDRITIS DISSECANS

The etiology of osteochondritis dissecans has given rise to much speculation, and in recent years the world literature has contained an increasing number of case reports bringing to light previously unknown features of the condition. Indeed the manifestations reported differ so widely from the previously accepted conception of the classical lesion that it has become more and more apparent that widely differing cases are concerned, and that the term osteochondritis dissecans merely serves to indicate a broad similarity between one case and another. No hard and fast criteria can be laid down as being necessary to justify the diagnosis, but it is usually accepted that the diagnosis can be properly applied to any case in which a portion of the articular surface, not necessarily containing bone, separates or attempts to separate by a process of gradual and “dissecting” extrusion.

There can be no doubt that in some cases—particularly in the knee and ankle—trauma, athletic or occupational over-use, or similar mechanical factors play an important part and may indeed be the sole cause of the condition. Even in these cases it is doubtful whether the lesion can ever be properly described as a simple osteochondral fracture. The fact that the lesion invariably shows itself on a convex articular surface and never on a concave one, although it remains unexplained, is also a clear indication that some mechanical factor must underlie the development of the lesion. Even when a constitutional factor is known to exist, as in bilateral affection of the elbow, it is always the dominant right arm that is more severely affected. However, mechanical factors alone are quite insufficient to explain the occurrence of multiple and often symmetrical lesions, familial and hereditary cases, and the association with dwarfing and endocrine disorder. White, in this number of the *Journal*, records three patients in whom the multiple lesions of osteochondritis dissecans were present in association with shortness of stature and fairly convincing evidence of endocrine upset. Local mechanical factors may determine the joints affected and the sites of the lesions, but the constitutional factor is obviously predominant. It is also interesting that in White's case there appears to be an association with osteochondritis of the spine which has also been noted by previous authors.

It is interesting to note the age incidence in different cases. Affection of one or both hip joints may occur in a young child. The lesion is radiographically identical with those seen later in life. There is a strong tendency to spontaneous healing of the lesion with conservative treatment, or even without treatment. It is also noticeable that when the constitutional factor seems to be preponderant, as shown by multiple joint involvement, symmetrical involvement of the knee caps and familial cases, the age incidence tends to be lower than in cases in which a single joint is involved. There appears to be a stronger case for the operation

of some constitutional factor in affection of the elbow or elbows, and the age incidence is much lower, the onset being almost invariably before growth is complete. By contrast, osteochondritis dissecans affecting the knee joint commonly arises after growth is complete and without any evidence of any constitutional cause.

With regard to treatment, there has been an increasing tendency towards a more conservative approach than was once popular. If the lesion can heal spontaneously in a young child, why should our treatment not be directed towards inducing healing of the lesion in older children and young adults by surgical or non-surgical means? In the past it has too often been accepted that separation of the detached fragment or fragments is inevitable, and surgeons have anticipated detachment of the loose fragment by carving it out, even at a stage when the overlying articular cartilage, although softened, remains unbroken. This sort of surgical excision is only justified if it is clear that there can be no hope of repair. There are occasions when the existence of a crater on the joint surface is less likely to lead to symptoms and subsequent arthritis than the persistence of irreparable detachment and fragmentation of cartilage and bone. The recommendation of Guilleminet and Barbier that excision of the lesion in the hip is the treatment of choice may be appropriate to the cases they describe, but it would be quite inappropriate in young children, and in many cases in older children and adults in whom symptoms are minimal or absent.

The practice of multiple drilling of a partly detached fragment has been employed for many years, although not much publicised. Used in the appropriate case it can lead to radiological evidence of healing of the lesion. An alternative procedure is to use a small peg bone-graft, countersunk in a small hole in the articular surface. Smillie has now, with great success, carried conservative surgical treatment a stage further by replacing in the crater a loose body which has already become detached and is lying free in some other part of the joint. It is doubtful whether such a policy is really necessary when the lesion lies in the so-called classical site on the medial condyle close to the intercondylar notch, but in other cases, which are just as common, in which the crater lies on an important weight-bearing surface of one or other femoral condyle, restoration of the articular surface is highly desirable and Smillie has shown that it can be achieved. It will, of course, take some time before the limitations of the method can be defined, but in the meantime Smillie's contribution serves to remind us of the possibilities of conservative surgery.

His suggestion that an extension of the principle could lead to the use of homografts of bone and articular cartilage is an interesting speculation, but it would be premature to consider this as a practical surgical possibility until the possibility of survival of such a graft has been proved experimentally.

NORMAN ROBERTS.