SPONDYLOLISTHESIS. A Study of 53 Cases Treated by Spine Fusion and 32 Cases Treated by Laminectomy. By L. E. LAURENT. *Acta Orthopaedica Scandinavica*. Supplement XXXV. From the Orthopaedic Hospital of the Invalid Foundation, Helsingfors, Head: Docent Anders Langenskiöld, M.D. $9\frac{3}{4} \times 6\frac{1}{4}$ in. Pp. 45, with 12 figures and 15 tables. 1958. Copenhagen: Ejnar Munksgaard Limited. Price Da. kr. 15.

This excellent little monograph from Finland does not pretend to be exhaustive but does provide a valuable review of the neurological changes and the surgical treatment. It is extraordinary that such a small subject should have attracted so much interest during the past forty years. Probably this is because of the etiological obscurity, the severe deformity which may result, and the surgical problems raised by attempts to reduce the deformity or to provide stability for an obviously weak section of the spine. During these years we have learnt that the initial lesion occurs during early childhood, that severe displacements may follow during adolescence, and that reduction more or less complete can sometimes be achieved; but it is extremely difficult, if not impossible, to retain this, nor does posterior bone grafting in adolescence provide any assurance against later further displacement. The writer of this paper reports upon seventy-seven of the patients who were treated surgically. Of these seventy-seven, forty-four were subjected to posterior spinal fusion, seven to anterior fusion and twenty-six to laminectomy without fusion. The results of fusion were better than those of laminectomy without fusion. The extent of slipping did not influence the results. The indication for operative treatment is persistent pain that is not relieved by conservative treatment. In adolescents whereolisthesis appears likely to lead to total displacement fusion should also be performed.—Norman CAPENER.

INJURIES AND SURGICAL DISEASES OF THE ISCHIUM. By Henry MILCH, M.D., Attending Orthopaedic Surgeon, Hospital for Joint Diseases, New York. $10\frac{1}{4} \times 7\frac{1}{4}$ in. Pp. x+163, with 106 figures. 1958. Index. New York: A Hoeber-Harper Book. London: Pitman Medical Publishing Co. Ltd. Price £4, 4s.

Any book solely devoted to a small specialised subject must inevitably stress points which can have very little practical value. This monograph is no exception. It gives an exhaustive account of the anatomy, injuries and diseases of the ischium. For example, the fact that "a fissure fracture may sometimes be overlooked" can hardly be of even academic interest. Similarly, the twelve pages devoted to the rather rare condition of ischial apophysiolyis prove to be tedious reading. There are over a hundred illustrations; but unfortunately many of the reproductions of radiographs are of very poor quality and this makes it very difficult to appreciate the points being shown.

Much of the work is a review of the literature and there is a full bibliography at the end of each chapter. This may prove of value to anyone who is engaged in research on the same subject. The best part of the book comes at the end, where there is an excellent chapter on the surgery of the ischium. The illustrations of the operation for excision of the ischium are particularly good. This part should be read by anyone who is contemplating excising this bone for tumour or other disease.—J. N. WILSON.

FRACTURE OF THE OS CALCIS AND ITS TREATMENT. II. A Contribution to the discussion on the treatment of calcaneus fracture based on an analysis of a ten-year material treated by closed reduction and traction, from Sentralsykehuset i Trondheim. By Arne ARNESEN. $9\frac{1}{2} \times 7$ in. Pp. 70, with 60 radiographs, 54 figures, 2 photographs and 21 tables. 1958. Stockholm: Acta Chirurgica Scandinavica, Supplement 234. Price Sw. kr. 10.

This monograph on the treatment of fractures of the calcaneum by Arnesen of Trondheim deals mainly with a detailed study of fifty-eight patients with this troublesome injury. The usual Norwegian thoroughness and attention to detail are pleasant features, and, although the English translation is somewhat quaint, the meaning is clear enough. The author is obviously honest in assessing his results and does not claim to have mastered the tongue-type or the severe stamp-type of deformity. He does claim, and with good reason, that divergent traction through the metatarsals and through the tuberosity of the calcaneum, plus manipulation, can wholly restore the tuber-joint angle, and lead to good results in 90 per cent of the depression type of fractures. The author described this method of treatment in 1939; in the present work he describes several changes in technique which he thinks have been responsible for an improvement in results. Traction is maintained for six weeks, because the ordinary calcaneal fracture is stable at the end of this time; plaster fixation has been given up; active exercises are considered essential, and are started as soon as traction has been applied. This is a stimulating and valuable contribution towards the solution of this difficult problem—the adequate treatment of fracture of the calcaneum.—L. W. PLEWES.