The Journal of Bone and Joint Surgery

EDITORIALS AND ANNOTATIONS

CONGENITAL DISLOCATION OF THE HIP

In 1953 Muller and Seddon reported the late results of treating congenital dislocation of the hip by the then best available methods: they were far from satisfactory. Somerville (1967), by his recognition of the importance of the inverted limbus, has remarkably improved the results of dislocations diagnosed weeks or months after birth. The result relates directly to the age at diagnosis. These patients usually require at least one operation and several weeks in hospital.

In this Journal in 1962 von Rosen and Barlow separately reported large series of neonatal examinations for congenital dislocation of the hip. It seemed from these two papers that dislocation could easily be detected at birth by clinical examination and treated by a simple splint. It seemed possible that surgery for this condition and the late serious sequelae would disappear. In some infants, the capsule being lax, the hip would dislocate if extended; if the dislocation persisted secondary changes gave the classical features of congenital dislocation of the hip. The splint prevented this.

In 1968 Smaill of New Zealand, and Emnéus, von Rosen, Hierton and James from Sweden reported further in this Journal. Though there were cases of missed dislocations and unsuccessful treatment by splintage the emphasis was on a truly dramatic change in the diagnosis and management of this condition.

The four papers in this issue are based on several years' use of the Ortolani and Barlow tests in the newborn and splintage for abnormal hips. In Edinburgh Mitchell works under favourable circumstances, a geographically closed community, an accessible and large obstetric unit and the virtual certainty, since he is the only orthopaedic surgeon treating congenital dislocation of the hip, that all missed cases and failures become known. By Mitchell's definition a luxation is present when a hip is demonstrably dislocatable by the Ortolani or Barlow test: all are splinted. An unstable hip is one in which a clunk has been briefly present: these hips are not splinted unless radiographic changes are present and none has later dislocated. In the years 1961-68, of 31,961 infants examined at birth only four were seen later with missed dislocation—one in 8,000 infants. During this period 100 luxations were treated by splints and ninety-two have normal hips. Eight failed to reduce or to be held reduced. Mitchell would now state that this is due to an infolded limbus. Timely excision of this repeats the excellent results reported by Somerville. Therefore Mitchell implies that of those diagnosed at birth only 8 per cent need operation and in these the results should be good. There remains the problem of one missed case in 8,000.

MacKenzie's report from Aberdeen (with the same population) confirms the Edinburgh report but shows curious differences. Perhaps his greatest achievement is to organise such a

very effective examination of infants over the north and north-west of Scotland, a sparsely inhabited area with many isolated communities. Some seventy dislocations were missed, 1·12 in 1,000 births (rarely in the peripheral centres staffed by general practitioners), almost as high as the incidence of congenital dislocation of the hip reported in Britain. The number of hips operated upon was also much greater. The number of abnormal hips, 1,671 in nearly 80,000 births, is three times greater than in Edinburgh. Is there more congenital dislocation in this area? It is possible genetically in North-East Scotland, for many inhabitants are Scandinavian in origin. Are MacKenzie's criteria different, particularly his group of "stiff hips"?

In contrast, from Northern Ireland, a widely scattered community with 30,000 births annually, Williamson reports that the referral of established cases of congenital dislocation has not fallen significantly and that there is a considerable problem from unsuccessfully splinted cases. The newborn are examined by a varied group of doctors. Two units carrying out intensive studies show quite remarkable differences of incidence of diagnosed and missed cases. It is a timely warning of the practical difficulties of examining all infants in a community.

Wilkinson has had similar difficulties. He emphasises the importance of breech births which yield a high incidence of congenital dislocation, perhaps because the legs are extended *in utero*—if so, why should this increase the number of hip dislocations?

We must accept that many babies are splinted unnecessarily—fifteen to twenty times the expected incidence of dislocation (Hierton and James 1968). We must ask how soon, or late, to splint; is three weeks too late? Some centres, such as Belfast, Aberdeen and even New Zealand, show that missed dislocations approximate to the expected incidence of congenital dislocation. Are persistent dislocations missed at birth as suggested by Owen (1968). Are we treating those hips which would cure themselves and not the real problem? Difficulties certainly are due to inexperienced examination of the newborn and the problems of organisation. Infants born by the breech or with a family history of hip dislocation need particular care. Perhaps all infants require successive examinations of the hips in the first one or two years of life.

Aberdeen has three times as many abnormal hips as Edinburgh: is this true or are criteria different? Finlay, Maudsley and Busfield (1967) in a paper most relevant to this problem suggested criteria and a terminology. Simplified, it might help to resolve our present uncertainties which stem more from difference of method and terminology than from a true failure of neonatal diagnosis and treatment of congenital dislocation.

There are suggestions that congenital dislocation may occur either because ligaments are lax or because the acetabular roof is ill formed, both anomalies being genetically influenced (Wynne-Davies 1970). To equate this latter as the reason for "missed" hips is attractive but as yet without supporting evidence (Walker 1971, Owen 1971, Corkery 1971).

With the excellent figures from Swedish authors, from Smaill and from Mitchell, it is difficult not to accept that neonatal examination carried out by experienced orthopaedic surgeons can, under ideal circumstances, almost solve the problem of congenital dislocation. If the few failures of splinting are, without delay, treated by appropriate surgery the results have indeed been shown, under ideal circumstances, to be excellent. J. I. P. JAMES.

REFERENCES

- BARLOW, T. G. (1962): Early Diagnosis and Treatment of Congenital Dislocation of the Hip. *Journal of Bone and Joint Surgery*, **44-B**, 292.
- CORKERY, P. H. (1971): Congenital Hip Dislocation. *British Medical Journal*, **3**, 371.
- EMNÉUS, H. (1968): A Note on the Ortolani-von Rosen-Palmén Treatment of Congenital Dislocation of the Hip. *Journal of Bone and Joint Surgery*, **50-B**, 537.
- FINLAY, H. V. L., MAUDSLEY, R. H., and BUSFIELD, P. I. (1967): Dislocatable Hip and Dislocated Hip in the Newborn Infant. *British Medical Journal*, **4**, 377.

- HIERTONN, T., and JAMES, U. (1968): Congenital Dislocation of the Hip. *Journal of Bone and Joint Surgery*, **50-B**, 542.
- MULLER, G. M., and SEDDON, H. J. (1953): Late Results of Treatment of Congenital Dislocation of the Hip. *Journal of Bone and Joint Surgery*, **35-B**, 342.
- OWEN, R. (1968): Early Diagnosis of the Congenitally Unstable Hip. *Journal of Bone and Joint Surgery*, **50-B**, 453.
- OWEN, R. (1971): Congenital Hip Dislocation. *British Medical Journal*, **3**, 370.
- ROSEN, S. von (1962): Diagnosis and Treatment of Congenital Dislocation of the Hip Joint in the New-born. *Journal of Bone and Joint Surgery*, **44-B**, 284.
- ROSEN, S. von (1968): Further Experience with Congenital Dislocation of the Hip in the Newborn. *Journal of Bone and Joint Surgery*, **50-B**, 538.
- SMAILL, G. B. (1968): Congenital Dislocation of the Hip in the Newborn. *Journal of Bone and Joint Surgery*, **50-B**, 524.
- SOMERVILLE, E. W. (1967): Results of Treatment of 100 Congenitally Dislocated Hips. *Journal of Bone and Joint Surgery*, **49-B**, 258.
- WALKER, G. (1971): Problems in the Early Recognition of Congenital Hip Dislocation. *British Medical Journal*, **3**, 147.
- WYNNE-DAVIES, R. (1970): Acetabular Dysplasia and Familial Joint Laxity: Two Etiological Factors in Congenital Dislocation of the Hip. *Journal of Bone and Joint Surgery*, **52-B**, 704.