Table I. The main features distinguishing reflex sympathetic dystrophy (RSD) and bone-marrow oedema syndrome (BMES)

	RSD	BMES
Aetiology and gender		
Previous trauma	Frequent (80%)	Uncommon
Risk of AVN	Occasional	Frequent (70%)
Female:male	3:1	1:2
Localisation and symptoms		
Joint prevalence	Hands and feet	Hip
Migratory or recurrent	Uncommon	Frequent (15% to 40%)
Onset and course	Chronic and continuous	Acute and progressive
Quality of pain	Burning with muscle spasm	Mechanical (on weight-bearing)
Trophic and vasomotor signs	Frequent (80%)	Uncommon
Permanent disability	Moderately often (20%)	Uncommon
Imaging		
Loss of radiodensity	Patchy or spotty	More homogeneous
Bone scans (increased uptake)	Diffuse (often periarticular)	More local
MRI		
Signal characteristics	Patchy	More homogeneous
Joint effusion	Moderately often (30%)	Frequent (90%)
Location	All articulating bones	Usually only one bone
Soft-tissue involvement	Frequent (90%)	Unusual (10%)
Functional exploration		
Bone-marrow pressure, stress test	Pathologically high in both	
Venography	Reflux and stasis in both	
Marrow changes	BME, marrow necrosis, fibrovascular regeneration	
Trabecular bone changes	New bone formation, bone necrosis, no osteoporosis	

RSD (Schimmerl et al 1991; Hofmann et al 1993) has identified significant differences between BMES and RSD (Table I).

In cases of ischaemia, there appears to be a common pathway in bone: bone-marrow oedema seems to be a uniform pattern of reaction in bone and its vascular supply to various external or internal disturbances (Rutishauser, Rohner and Held 1960). It is therefore not surprising that no significant differences are found in the radiological appearance or at the functional exploration of bone. Despite this we agree with Lakhanpal et al (1987) that BMES is a syndrome of monoarticular or oligoarticular pain in young or middle-aged people with significant differences from RSD. In contrast to most other authors, however, we believe that BMES is an early reversible form of avascular necrosis (AVN) and not a distinct disease.

S. HOFMANN, MD
Danube Hospital
Vienna, Austria.
H. PLENK, Jr, MD
University of Vienna
Vienna, Austria.

Fialka V. The diagnosis of reflex sympathetic dystrophy. *Eur J Phys Med Rehabil* 1992; 2:40-4.

Hofmann S, Kramer J, Leder K, et al. Correlation of MRI and histomorphological findings in bone marrow oedema syndrome of the hip. *Eur Radiol* 1993; 3:408-12.

Hofmann S, Kramer J, Leder K, Plenk H Jr, Engel A. Bone marrow edema syndrome (BMES) of the hip: clinical and MR-imaging follow-up after core decompression treatment. *Orthop Trans AAOS* 61th annual meeting, New Orleans, 1994.

Kramer J, Hofmann S, Engel A, Leder K, Neuhold A, Imhof H. Hüftkopfnekrose und Knochenmarksödemsyndrom in der Schwangerschaft. *Fortschr Röntgenstr* 1993; 159:126-31.

Lakhanpal S, Ginsburg W, Luthra H, Hunder G. Transient regional osteoporosis: a study of 56 cases and review of the literature. *Ann Int Med* 1987; 106:444-50.

Rutishauser E, Rohner A, Held D. Experimentelle Untersuchungen über die Wirkung der Ischämie auf den Knochen und das Mark. *Virchows Arch Path Anat* 1960; 333:101-18.

Schimmerl S, Schurawitzky H, Imhof H, et al. Morbus Sudeck: MRT als neues diagnostisches Verfahren. *Fortschr Röntgenstr* 1991; 154:601-4.

Veldman P, Reynen H, Arntz I, Goris J. Signs and symptoms of reflex sympathetic dystrophy: prospective study of 829 patients. *Lancet* 1993; 342:1012-6.

AETIOLOGY OF CLUB FOOT

Sir,

The article in the March 1994 issue entitled 'Posterolateral release for resistant club foot' by Hudson and Catterall (1994; 76-B:281-4) raises the question of the role of the subtalar joint in the pathogenesis of club foot. Most operative approaches provide subtalar release as the primary treatment, and therefore the good results of posterolateral release suggest that internal talar rotation, not subtalar varus, is the primary pathological problem.

This has previously been stressed by Goldner (1969) who found that the primary deformity was medial and volar angulation of the neck of the talus. He also described the 'lateral impingement syndrome', for which a posterolateral release was used in addition to medial derotation involving the deltoid and medial capsular structures of the ankle, for severely contracted feet.

The good results of posterolateral release again emphasise that the subtalar joint need not be released to obtain such results in many cases. If the deformity is more severe, the combination of primary medial release of the talus at ankle level and posterolateral release is indicated.

H. P. AIDEM, MD
Phoenix Orthopaedic Group
Phoenix,
Arizona, USA.

Goldner JL. Congenital talipes equinovarus = fifteen years of surgical treatment. *Curr Pract Orthop Surg* 1969; 4:61-123.

Hudson I, Catterall A. Posterolateral release for resistant club foot. *J Bone Joint Surg [Br]* 1994; 76-B:281-4.