Correspondence

We welcome letters to the Editor concerning articles which have recently been published. Such letters will be subject to the usual stages of selection and editing; where appropriate the authors of the original article will be offered the opportunity to reply.

Letters should normally be under 300 words in length, double-spaced throughout, signed by all authors and fully referenced. The edited version will be returned for approval before publication.

Anatomy of the medial femoral circumflex artery and its surgical implications

Sir,

I write with reference to the paper by Gautier et al¹ entitled 'Anatomy of the medial femoral circumflex artery and its surgical implications' in the July 2000 issue.

The authors have attempted to focus attention on the importance of knowledge of the blood supply of the upper end of the femur in surgical practice.

The paper is based on anatomical dissections of 12 fresh cadavers and it is supported by 52 references made in *en bloc* form. Many of the statements which the authors make, however, are not supported by the work of other researchers, for example, "The deep branch of the MFCA gives rise to two to four superior retinacular vessels and occasionally, to inferior retinacular vessels." The work of Trueta and Harrison² was shown to be flawed many years ago by Lagrange and Dunoyer³ whose work is not quoted in the references, and by Crock⁴ and Chung⁵.

The fact that multiple retinacular vessels have not been demonstrated by the present authors may indicate that they were not fully injected during the preparation of the specimens, rather than that they are absent.

Another statement made is: "Despite previous descriptions,^{9,12,14,17,18,20-22,24,49} we found no anastomotic branch surrounding the neck of the femur . . ." The ten cited references are merely listed, and several are outstanding anatomical works which disprove this claim.

Unfortunately, this paper simply perpetuates fallacies about the blood supply of the upper end of the femur since it is based on the use of inadequate anatomical techniques. Also, the five authors thank six co-workers for technical assistance and friendship, raising serious questions about multiauthorship. For example, who did the dissections, and who accepts the academic responsibility for conclusions drawn from the work?

H. V. CROCK, AO, MS, MD, FRCS, FRACS, FRCS Ed (Hon)
Spinal Disorders Unit
Cromwell Hospital, London, UK.

1. Gautier E, Ganz K, Krügel N, Gill T, Ganz R. Anatomy of the medial femoral circumflex artery and its surgical implications. *J Bone Joint Surg [Br]* 2000;82-B:679-83.

2. Trueta J, Harrison MHN. The normal vascular anatomy of the femoral head in adult man. *J Bone Joint Surg [Br]* 1953;35-B:442-61.

3. Lagrange J, Dunoyer J. La vascularisation de la tête fémorale de l'enfant. *Rev Chir Orthop Reparatrice Appar Mot* 1962;48:123-37.

4. Crock HV. A revision of the anatomy of the arteries supplying the upper end of the human femur. *J Anat Lond* 1965;99:77-88.

5. Chung SMK. The arterial supply of the developing proximal end of the human femur. *J Bone Joint Surg [Am]* 1976;58-A:961-70.

Author's reply:

Sir,

The main aim of our paper was to advance knowledge of surgical anatomy in its application to safe and reproducible surgery of the hip.

In his letter, Mr Crock criticises a perceived deficiency in the accuracy of our study because of inadequate vascular perfusion using rubber latex. He has based this on the (wrong) assumption that we were dealing with the entire vessel distribution supplying the proximal end of the femur. It was our intention, however, to describe the applied surgical anatomy of the ramus profundus of the medial femoral circumflex artery along the external rotators. This is of practical importance and can be seen by the surgeon in his daily work. The study was initiated because most atlases of anatomy and textbooks of approaches in use today give only a minimum amount of information on the anatomy of the medial femoral circumflex artery which is useful surgically. This is especially true of the course of the vessel along the external rotators.

Perfusion with latex rubber has a long and successful tradition in the anatomy laboratory in Paris where we have performed several studies, including the present one.¹⁻³ It can fill vessels with a diameter of surgical interest of 0.5 mm and greater. The same technique was used for data in the books of Tubiana, McCullough and Masquelet⁴ and Masquelet, McCullough and Tubiana⁵ on surgical approaches to the extremities. In our present study, we used only specimens in which excellent perfusion could be demonstrated up to the entrance of the vessels into the femoral head. We therefore believe that we have largely excluded any significant vascular or technical variations. Moreover, if arteries were present on surfaces and failed to fill, they would certainly not be as patent or extensive as the adjacent vessels, which filled very well.

In his further critique, Mr Crock expounds on the vascular anatomy of the proximal femur in childhood with his reference to Lagrange and Dunoyer and Chung. Our study examined the vascular anatomy in the adult exclusively. Although this is not mentioned specifically, it is alluded to within the text. There is little doubt that the vascular distribution in early childhood and in the adult have some differences. One example is the regression of the lateral femoral circumflex artery with age, a phenomenon which was mentioned by Ogden⁶ and which explains why there is no substantial peripheral anastomosis between the medial and lateral femoral circumflex artery in adulthood. This is also stated by Gillot, Sakka and Frot⁷ in their detailed anatomical study.

The findings of our study are constantly confirmed in our daily surgical practice, including routine dislocation of the hip at operation. They have been reinforced recently by dynamic perfusion profiles of the femoral head during surgical dislocation, using intraosseous laser-Doppler flowmetry. We thus feel that Mr Crock's reproach that we simply perpetuate fallacies about the

blood supply of the upper end of the femur, is not justified. This paper has information important for hip surgery which is not available from other sources.

E. GAUTIER, MD
Hôpital Cantonal
Fribourg, Switzerland.

1. Ballmer FT, Masquelet AC. The reversed-flow medio-distal fascio-cutaneous island thigh flap: anatomic basis and clinical applications. *Surg Radiol Anat* 1998;20:311-6.
2. Ballmer FT, Hertel R, Nötzli HP, Masquelet AC. The medial malleolar network: a constant vascular base of the distally based saphenous neurocutaneous island flap. *Surg Radiol Anat* 1999;21:297-303.
3. Gerber C, Schneederger AG, Vinh TS. The arterial vascularization of the humeral head. *J Bone Joint Surg [Am]* 1990;72-A:1486-94.
4. Tubiana R, McCullough CJ, Masquelet AC. *Voies d'abord chirurgicales du membre supérieur*. Paris: Masson, 1992.
5. Masquelet AC, McCullough CJ, Tubiana R. *An atlas of surgical exposure of the lower extremity*. London: Martin Dunitz Ltd, 1993.
6. Ogden JA. Changing patterns of proximal femoral vascularity. *J Bone Joint Surg [Am]* 1974;56-A:941-50.
7. Gillot C, Sakka M, Frot B. The internal circumflex artery of the thigh. *Arch Anat Pathol* 1971;19:397-408.

Nonunion of the femoral diaphysis

Sir,

I read with interest the article in the July 2000 issue by Giannoudis et al¹ entitled 'Nonunion of the femoral diaphysis.'

The main finding of their retrospective review of 377 patients with fractures of the femoral diaphysis was the association between delayed and nonunion and the use of non-steroidal anti-inflammatory drugs (NSAIDs). This was interpreted as being causative because the authors state that they now exclude the use of NSAIDs in these patients.

Some animal studies support this hypothesis,²⁻⁶ while others do not.^{7,8} There are few human studies on the effect of NSAIDs on the healing of fractures. Butcher and Marsh's abstract⁹ quoted in this paper as supporting evidence, was also a retrospective study. Davis and Ackroyd¹⁰ found no effect of flurbiprofen on bone healing compared with placebo in a double-blind prospective trial of Colles' fractures.

An alternative interpretation of the authors' findings is that patients with delayed nonunion have pain, and this results in usage of NSAIDs.

Rather than excluding use of NSAIDs in these patients, we think that a more measured response to the findings of this paper would be a randomised controlled trial. NSAIDs are popular analgesics with anaesthetists and general practitioners and patients often take NSAIDs for a variety of medical conditions. The use of aspirin in orthopaedic patients may increase because aspirin has been shown to reduce clinical deep-vein thrombosis and the rate of pulmonary embolism after fracture of the neck of the femur.¹¹ We need to know the risk of nonunion with NSAIDs, so that we can weigh up the benefits and risks when making decisions about their use for our patients with fractures.

If we interpret the odds ratio of Giannoudis et al¹ for the use of NSAIDs conservatively, and assume the risk of nonunion with NSAIDs to be fivefold, to have an 80% chance of demonstrating a fivefold increase in the rate of nonunion at a significance level of $p = 0.05$, would require a trial of 150 patients.

R. K. PRATT, FRCS
D. J. KRAMER, FCS(SA) Orth
North Tyneside General Hospital
Newcastle upon Tyne, UK.

1. Giannoudis PV, MacDonald DA, Matthews SJ, et al. Nonunion of the femoral diaphysis: the influence of reaming and non-steroidal anti-inflammatory drugs. *J Bone Joint Surg [Br]* 2000;82-B:655-8.
2. Dimar JR II, Ante WA, Zhang YP, Glassman SD. The effects of nonsteroidal anti-inflammatory drugs on posterior spinal fusions in the rat. *Spine* 1996;21:1870-6.
3. Engesaeter LB, Sudmann B, Sudmann E. Fracture healing in rats inhibited by locally administered indomethacin. *Acta Orthop Scand* 1992;63:330-3.
4. Høgevoid HE, Groggaard B, Reikeras O. Effects of short-term treatment with corticosteroids and indomethacin on bone healing: a mechanical study of osteotomies in rats. *Acta Orthop Scand* 1992;63:607-11.
5. Lindholm TS, Tornkvist H. Inhibitory effect on bone formation and calcification exerted by the anti-inflammatory drug ibuprofen: an experimental study on adult rat with fracture. *Scand J Rheumatol* 1981;10:38-42.
6. Allen HL, Wase A, Bear WT. Indomethacin and aspirin: effect of nonsteroidal anti-inflammatory agents on the rate of fracture repair in the rat. *Acta Orthop Scand* 1980;51:595-600.
7. Reikeras O, Engebretsen L. Effects of ketoralac tromethamine and indomethacin on primary and secondary bone healing: an experimental study in rats. *Arch Orthop Trauma Surg* 1998;118:50-2.
8. Huo MH, Troiano NW, Pelker RR, Gundberg CM, Friedlaender GE. The influence of ibuprofen on fracture repair: biomechanical, biochemical, histologic and histomorphometric parameters in rats. *J Orthop Res* 1991;9:383-90.
9. Butcher CK, Marsh DR. Non-steroidal anti-inflammatory drugs delay tibial fracture union. *Injury* 1996;27:375.
10. Davis TRC, Ackroyd CE. Non-steroidal anti-inflammatory agents in the management of Colles' fractures. *Br J Clin Pract* 1988;42:184-9.
11. Rodgers A, MacMahon S, Collins R, Prentice C. Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: pulmonary embolism prevention (PEP) trial. *Lancet* 2000;355:1295-302.

Author's reply:

Sir,

We agree that the most powerful way to assess the effect of NSAIDs is by a proper randomised clinical trial and look forward to this being performed. The experience in trauma surgery, however, and in a wide variety of clinical situations, is that bone healing is restricted by NSAIDs. Bone healing is multifactorial and this trial was set up to look at the effect of reamed against unreamed nails because of our previous report describing delay in healing and, more recently, the increased rate of nonunion with unreamed nails noted by the Canadian group.¹ The effect of the NSAIDs was so marked, however, that it effectively swamped the other variables in our groups. We have had our data reassessed by several statisticians and do believe that this is a real effect. The clinical impression after removal of NSAIDs from our practice has been the disappearance of delayed and occasionally, poor healing in fractures of the femoral diaphysis.

R. M. SMITH, MD, FRCS
Trauma Service, St James's University Hospital
Leeds, UK.

1. Powell JN, DeGroot R, Kreder HJ, et al. A prospective, randomised trial comparing reamed and unreamed intramedullary nailing: an analysis of rates of union. Orthopaedic Trauma Assoc, October 1999.

Sir,

We read with interest the article by Giannoudis et al¹ in the July 2000 issue entitled 'Nonunion of the femoral diaphysis.'

We have some concerns regarding the outcome measures used to define union. The clinical and radiological methods used to define healing are inaccurate.^{2,3} The radiographs were not standardised; this adds further to the inaccuracies of a method relying on the quantification of callus to define union. The need for surgical intervention is dependent on this definition as is the separation into the two study groups. Nonunion was found to occur with a mean time to diagnosis of 11.5 months (7 to 18) in the study group, and 6 months (3 to 10) in the control group. This overlap questions the methods of defining union.

We note the effects of varying factors on union, and question the conclusions drawn since inaccurate outcome measures were used. The deleterious effects of non-steroidal anti-inflammatory drugs (NSAIDs) on healing in the nonunion group are discussed, although the timing of the administration of NSAIDs is not recorded. Patients with nonunion may continue to take analgesics as their fractures are still painful, and they bear weight later at 7.8 weeks compared with 4 weeks. Although this is stated in the discussion, the administration of NSAIDs for their analgesic effect seems to have been ignored. It is therefore difficult to separate the 'cause and effect' of the administration of NSAIDs and nonunion.

Studies using inaccurate outcome measures should be interpreted with care. The study of the healing of fractures is no exception.

R. H. WADE, FRCS Ed
P. J. ROBERTS, FRCS
J. B. RICHARDSON, MD, FRCS
Robert Jones & Agnes Hunt Orthopaedic Hospital
Oswestry
Shropshire, UK.

1. Giannoudis PV, MacDonald DA, Matthews SJ, et al. Nonunion of the femoral diaphysis: the influence of reaming and non-steroidal anti-inflammatory drugs. *J Bone Joint Surg [Br]* 2000;82-B:655-8.
2. Hammer R, Hammerby S, Lindholm B. Accuracy of radiologic assessment of tibial shaft fractures in humans. *Clin Orthop* 1985;199:233-8.
3. Hammer R, Norrbom H. Evaluation of fracture stability: a mechanical simulator for assessment of clinical judgement. *Acta Orthop Scand* 1984;55:330-3.

Author's reply:

Sir,

I would like to thank Messrs Wade, Roberts and Richardson for their comments. I am sure that we would all agree that there are some limitations in retrospective reviews, and our study is no different from others. The paper, however, was reviewed by at least two expert statisticians who did feel that it was sound, although the limitations described are indeed there. I also note the difficulty in defining union. This is present in all papers which discuss the union of diaphyseal fractures in the presence of an implant, because all of the measurements of stiffness are useless. I suppose that the most important feature is that, after statistical review, it became clear that the effects of the NSAIDs swamped all of the other issues which we were trying to examine, and appeared to be the important factor. Clearly, a properly conducted, prospective, randomised control trial is required and I look forward to it being undertaken.

With regard to the second point, it seemed that the earlier NSAIDs were given the larger the effect. I agree that the patient with nonunion may continue to take analgesics for longer than others, but most of the NSAIDs were given early in the course of treatment before long-term pain from nonunion became a relevant feature. I agree that cause and effect in a retrospective study cannot be separated. However, we have identified what I believe

to be a true association between NSAIDs and nonunion, which requires further investigation.

R. M. SMITH, MD, FRCS
Trauma Service
St James's University Hospital
Leeds, UK.

Does 'canal clearance' affect neurological outcome after thoracolumbar burst fractures?

Sir,

We read with interest the review article in the July 2000 issue by Boerger et al¹ entitled 'Does 'canal clearance' affect neurological outcome after thoracolumbar burst fracture?' We were, however, uncertain of some of the arguments.

First, we found the comparison of results for a selection of patients after either operative or non-operative treatment, questionable. The mean improvements in Frankel grade are stated, and it is noted that "the trend is for better results to be obtained in the non-surgical group". The calculation of a mean would appear to be inappropriate, since it is based on a grading system lacking a numerical measure of a continuous variable.² Furthermore, the discussion of selection bias is inadequate. Differences between the groups could mean that however they are compared, it would be difficult to base valid conclusions on the comparison.

Secondly, the contribution made by the discussion of complications, which were "reported in 75% of papers," is limited. Whether the incidence of complications in general is reassuringly low or worryingly high cannot be easily gauged. The note that one paper³ reports "17% of patients who were initially neurologically intact suffered deterioration after surgery" would appear to be a misquotation. In the paper referred to, neurological deterioration is described in 17% of patients in a group treated non-operatively, some of whom improved after subsequent operative intervention. These points detract from what comes across as "the case against 'canal clearance'". Surely caution is needed in interpreting masses of pooled information or 'evidence' concerning the value of a procedure. As implied in the review, patients should not be subjected to unjustified operations and attendant complications. Neither should selected cases be denied interventions that may be of net benefit. To what extent would the authors consider that further prospective study is required to answer fully the question that entitles their article?

P. T. CROSSMAN, FRCS
J. M. SCOTT, FRCS, Dip Math
Luton and Dunstable Hospital NHS Trust
Luton, UK.

1. Boerger TO, Limb D, Dickson RA. Does 'canal clearance' affect neurological outcome after thoracolumbar burst fractures? *J Bone Joint Surg [Br]* 2000;82-B:629-35.
2. Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. *Br Med J* 1983;286:1489-93.
3. Denis F, Armstrong GW, Searls K, Matta L. Acute thoracolumbar burst fractures in the absence of neurologic deficit: a comparison between operative and non-operative treatment. *Clin Orthop* 1984;189:142-9.

Authors' reply:

Sir,

We thank Messrs Crossman and Scott for their comments. The point is taken that a mean score is not entirely appropriate for

these data. The differences which we observed are not significant and an effect is not going to be proved whatever tests the observations are subject to. We simply wished to illustrate that the data available are poor, yet we are using these data in our day-to-day decisions on whether a patient needs surgery or not.

The correspondents agree that patients should not be subjected to unjustified operations and attendant complications. We would also agree that many patients benefit from surgery to stabilise the spine.

The suggestion that a prospective study could be helpful is never a bad one, although there ought to be considerable retrospective evidence in favour of the proposition before the prospective study addresses how beneficial the treatment is, and for what type of patient, in what circumstances. The fact that there is no evidence that patients benefit neurologically from canal clearance, even in the selected cases, is for the very simple reason that the damage is done at the moment of injury.

R. A. DICKSON, MA, ChM, FRCS, DSc
D. LIMB, BSc, FRCS Ed Orth
St James's University Hospital
Leeds, UK.

Thromboprophylaxis – which treatment for which patients?

Sir,

I read with much interest the Editorial by Prentice¹ in the May 2000 issue entitled 'Thromboprophylaxis – which treatment for which patients?', and various articles in the journal on the use of thromboprophylaxis after total hip and knee replacements.

Although there is controversy in the literature about the risks and benefits of thromboprophylaxis after TKR and THR, most orthopaedic surgeons use some kind of thromboprophylaxis.² What was not stressed was the duration of its use. There is evidence that thrombosis and its complications can develop up to three months³ after hip or knee replacement, but the current practice is to use thromboprophylaxis for 7 to 10 days, the duration of the hospital stay. Surely those who believe in the efficacy of thromboprophylaxis should use it in some form for three months.

S. KAPOOR
Glasgow Royal Infirmary
Glasgow, UK.

1. Prentice CRM. Thromboprophylaxis – which treatment for which patient? *J Bone Joint Surg [Br]* 2000;82-B:483-5.
2. Gillespie W, Murray D, Gregg PJ, Warwick D. Risks and benefits of prophylaxis against venous thromboembolism in orthopaedic surgery. *J Bone Joint Surg [Br]* 2000;82-B:475-9.
3. Planes A, Vochelle N, Darmon JY, et al. Risk of deep-venous thrombosis after hospital discharge in patients having undergone total hip replacement: double-blind randomised comparison of enoxaparin versus placebo. *Lancet* 1996;348:224-8.

Author's reply:

Sir,

Dr Kapoor raises the important point of whether thromboprophylaxis after orthopaedic surgery should continue after discharge from hospital. I agree with him that symptomatic deep-vein thrombosis and pulmonary embolism, both fatal and non-fatal, may occur after the 7- to 10-day period, at which time thromboprophylaxis has usually been stopped. I do not believe, however, that the current evidence is strong enough to support the routine use of heparin on an outpatient basis after hospital discharge. The published studies have used the surrogate endpoints of routine venography and, in some cases, ventilation perfusion scanning, to assess the effect of subcutaneous heparin after discharge.¹⁻³ Although there is a reduction in these endpoints by heparin, it is still not certain whether they can be extrapolated into clinical benefit for the patient. In the only trial to use symptomatic venous thromboembolism as the outcome to assess low-molecular-weight heparin after hip or knee replacement, there was no convincing benefit for extended thromboprophylaxis.⁴ There is an urgent need to carry out large randomised controlled trials in which short and prolonged prophylaxis are compared using clinical endpoints. The only large study to evaluate this in a controlled manner is the Pulmonary Embolism Prevention (PEP) trial in which aspirin, 160 mg daily, started preoperatively and continued for 35 days postoperatively, was compared with placebo.⁵ This trial showed that mortality in the aspirin group continued to be reduced, compared with the placebo group, in the interval between discharge from hospital and 35 days, and that the protective effect of aspirin continued from week 2 until week 5 postoperatively. Whether extending thromboprophylaxis after five weeks can produce clinical benefits, however, is still not known.

C. R. M. PRENTICE, MD, FRCP
Division of Medicine, University of Leeds, The Martin Wing
The General Infirmary
Leeds, UK.

1. Planes A, Vochelle N, Darmon JY, et al. Risk of deep-venous thrombosis after hospital discharge in patients having undergone total hip replacement: double-blind randomised comparison of enoxaparin versus placebo. *Lancet* 1996;348:224-8.
2. Dahl OE, Andreassen G, Aspelin T, et al. Prolonged thromboprophylaxis following hip replacement surgery: results of a double-blind, prospective, randomised, placebo-controlled study with dalteparin. *Thromb Haemost* 1997;77:26-31.
3. Bergqvist D, Benoni G, Björngrell O, et al. Low-molecular-weight heparin (enoxaparin) as prophylaxis against venous thromboembolism after total hip replacement. *N Engl J Med* 1996;335:696-700.
4. Heit JA, Elliott CG, Trowbridge AA, et al. Ardeparin sodium for extended out-of-hospital prophylaxis against venous thromboembolism after total hip or knee replacement: a randomised, double-blind, placebo-controlled trial. *Ann Intern Med* 2000;132:853-61.
5. Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: pulmonary embolism prevention (PEP) trial. *Lancet* 2000;355:1295-302.