Correspondence

We welcome letters to the Editor concerning articles which have recently been published. Such letters will be subject to the usual stages of selection and editing; where appropriate the authors of the original article will be offered the opportunity to reply.

Letters should normally be under 300 words in length, double-spaced throughout, signed by all authors and fully referenced. The edited version will be returned for approval before publication.

Anatomical reduction of intra-articular fractures of the distal radius

Sir,

I read with interest the paper in the January 2000 issue by Mehta, Bain and Heptinstall,¹ entitled 'Anatomical reduction of intra-articular fractures of the distal radius: an arthroscopically-assisted approach', and congratulate the authors on their work to improve treatment of fractures of the distal radius.

I would like to comment, however, on the statement that: "There was a significant difference ($p < 0.05$) in the incidence of pain (VAS >2) for the patients with no step (18%), <1 mm step (38%) and >1 mm step (100%)." The 100% appears to refer to one patient in the group with >1 mm step in the articular surface of the distal radius on postoperative radiography. No statistical test can reasonably be applied to data containing only one patient in a group. Therefore the assertion that the results show a "significant difference" is invalid. The best conclusion that can be made from these results is that there is a trend suggesting that increasing articular steps are associated with more pain. I feel that it is important not to make conclusions which go beyond those which can be supported by the available results.

T. E. J. HEMS, MA, DM, FRCS, FRCS Ed(Orth)
Victoria Infirmary,
Glasgow, UK.

1. Mehta JA, Bain GI, Heptinstall RJ. Anatomical reduction of intra-articular fractures of the distal radius: an arthroscopically-assisted approach. *J Bone Joint Surg [Br]* 2000;82-B:79-86.

Meta-analysis of thromboembolic prophylaxis after total knee arthroplasty

Sir,

I read with interest the article entitled 'Meta-analysis of thromboembolic prophylaxis after total knee arthroplasty' in the August 2000 issue by Westrich et al.¹ The conclusions reached by the authors are in keeping with those of the Fifth ACCP Consensus Conference on Antithrombotic Therapy.²

The incidence of deep-vein thrombosis (DVT) in the intermittent pneumatic compression device (IPCD) group was found to

be the lowest. This result needs to be interpreted with some caution since there are few studies assessing the efficacy of IPCD: the combined patient numbers in the warfarin and low-molecular-weight heparin (LMWH) trials are more than twice the combined patient numbers in the IPCD studies. Consequently, more confidence can be placed in the results with warfarin or LMWH prophylaxis.

While we would support the use of IPCD for thromboprophylaxis in knee arthroplasty we are surprised that the authors advocate their use with aspirin, when their own meta-analysis and previous studies in knee surgery have shown aspirin to offer no significant advantage over a placebo in preventing DVT or pulmonary embolism. Until there is further clinical evidence to the contrary we would recommend that IPCD be combined with warfarin or LMWH, but not aspirin.

LMWH is described as containing "a specific tetrasaccharide which binds antithrombin III". The active binding site of both LMWH and unfractionated heparin is in fact a pentasaccharide sequence and is responsible in part for the anticoagulant mechanism of action of these drugs.

R. ALIKHAN, MD
A. T. COHEN, MD, FRACP
King's College Hospital,
London, UK.

1. Westrich GH, Haas SB, Mosca P, Peterson M. Meta-analysis of thromboembolic prophylaxis after total knee arthroplasty. *J Bone Joint Surg [Br]* 2000;82-B:795-800.
2. Clagett GP, Anderson FA Jr, Geerts W, et al. Prevention of venous thromboembolism. *Chest* 1998;114:531-60.
3. Choay J. Biologic studies on chemically synthesized pentasaccharide and tetrasaccharide fragments. *Semin Thromb Hemost* 1985;11:81-5.

Author's reply:

Sir,

We thank Dr Alikhan and Dr Cohen for their interest on our study and their recognition that our conclusions are in accordance with the Fifth ACCP Consensus Conference on Antithrombotic Therapy.

Meta-analysis is a useful technique in which multiple studies are combined to increase statistical power. In our particular study, we reviewed the literature between January 1980 and December 1997 and identified 122 articles. We used strict inclusion criteria and a scoring system so that only 23 studies were deemed to be appropriate for the meta-analysis. This left over 6000 patients in the study cohort, a large group of patients. We attempted to minimise any bias by using a scoring system and by setting the requirement of a strong methodology for the studies selected. I disagree with the statement that there were "fewer studies assessing the efficacy of IPCD", since seven of the 23 studies had at least one arm which included IPCD (see Table I).

We believe that IPCD is the best form of prophylaxis for thromboembolic disease after total knee arthroplasty. It may be combined with aspirin, low-molecular-weight heparin, or warfarin. There are no adequate data to support one combination of prophylaxis with IPCD over another. In our previous article published in 1996,¹ a significant decrease in the prevalence of DVT was shown when aspirin was used with pneumatic compression compared with aspirin alone: 27% v 59%, respectively. We

are at present performing further research to define the role of other types of prophylaxis used in conjunction with pneumatic compression.

G. H. WESTRICH, MD
Hospital for Special Surgery,
New York, USA.

1. Westrich GH, Sculco TP. Prophylaxis against deep venous thrombosis after total knee arthroplasty: pneumatic plantar compression and aspirin compared with aspirin alone. *J Bone Joint Surg [Am]* 1996;78-A:826-34.

Proximal femoral bone loss and increased rate of fracture with a proximally hydroxyapatite-coated femoral component

Sir,

I read with interest the article entitled 'Proximal femoral bone loss and increased rate of fracture with a proximally hydroxyapatite-coated femoral component' by Radl et al¹ in the November 2000 issue.

I have always stated emphatically that, for a hydroxyapatite-coated implant to become integrated, the conditions of contact, stillness and stressing must be fulfilled. The hydroxyapatite coating itself does not give initial stability, as claimed by the authors. It is the design of stem which must provide initial total immobility (primary fixation) so that bone ingrowth (secondary fixation) can occur.

This concept is expounded by Furlong and Osborn² and by the clinical results shown by McNally et al³ in which a success rate of 99% was achieved at a mean period of follow-up of ten years. Primary mechanical fixation provided by the cone, collar and lateral fin is essential in resisting subsidence and rotation. Clearly, the implant used by the authors had no such attributes and consequently was not initially stable. It failed, but it must be understood that this was not the fault of the hydroxyapatite coating.

R. FURLONG, FRCS
Furlong Research Foundation,
London, UK.

1. Radl R, Aigner C, Hungerford M, Pascher A, Windhager R. Proximal femoral bone loss and increased rate of fracture with a proximally hydroxyapatite-coated femoral component. *J Bone Joint Surg [Br]* 2000;82-B:1151-5.
2. Furlong R, Osborn JF. Fixation of hip prostheses by hydroxyapatite ceramic coatings. *J Bone Joint Surg [Br]* 1991;73-B:741-5.
3. McNally SA, Shepperd JA, Mann CV, Walczak JP. The results at nine to twelve years of the use of the hydroxyapatite-coated femoral stem. *J Bone Joint Surg [Br]* 2000;82-B:378-82.

Authors' reply:

Sir,

We thank Mr Furlong for his interest in our article. He is concerned because the high rate of failure of the cementless stem may be related to the hydroxyapatite coating. This has been shown to result in early and sound integration of the implant in several histomorphometric studies.^{1,2} The formation of new bone, which is enhanced by the hydroxyapatite, seems to support biological osseointegration of the prosthetic device. Bone remodeling, however, caused by load transfer, cannot be influenced by the coating. Despite the benefits of hydroxyapatite the loss of coating is described as a definite phenomenon, accelerated by loading.

Mr Furlong is correct that the hydroxyapatite coating itself can never be the major factor in prosthetic stability. The straight, collarless implant used in this retrospective study, showed early migration and distal load transfer. It was stressed by the authors that the design of the stem is responsible for the mechanical stability of the implant and that hydroxyapatite coating does not compensate for a poor design. To the best of our knowledge this study is the first to provide clinical evidence for this unsurprising phenomenon and thus underlines the theoretical considerations of Mr Furlong.

R. RADL, MD
R. WINDHAGER, MD
University of Graz, Austria.

1. Furlong RJ, Osborn JF. Fixation of hip prostheses by hydroxyapatite ceramic coatings. *J Bone Joint Surg [Br]* 1991;73-B:741-5.
2. Hardy DCR, Frayssinet P, Guilhem A, Lafontaine MA, Delinc PF. Bonding of hydroxyapatite-coated femoral prostheses: histopathology of specimens from four cases. *J Bone Joint Surg [Br]* 1991;73-B:732-40.
3. Tonino AJ, Therin M, Doyle CH. Hydroxyapatite-coated femoral stems: histology and histomorphometry around five components retrieved at postmortem. *J Bone Joint Surg [Br]* 1999;81-B:148-54.

The orthopaedic management of peripheral ischaemia in meningococcal septicaemia in children

Sir,

We read with interest the article in the April 2000 issue by Davies et al¹ entitled 'The orthopaedic management of peripheral ischaemia in meningococcal septicaemia in children', taking note particularly of the proposed protocol for treating such children.

Did the authors take into account the findings of our paper² in which we reviewed 12 infants and children suffering from peripheral limb ischaemia? We concluded that early surgical intervention with thrombectomy or embolectomy alone was usually unsuccessful in these difficult cases. We cited several other studies which recommended that postoperative intravenous infusion of heparin could help to reduce the progression of ischaemic damage in the limbs of these children.

This advice comes on the knowledge that in purpura fulminans, circulating immune complexes activate complement on the walls of small vessels leading to intimal damage and thrombosis. The use of heparin limits the degree of the thrombosis in the microcirculation, thus reducing the extent of the ischaemic damage.

M. J. FARRAR, FRCS Ed(Orth)
G. C. BENNETT, FRCS
N. I. L. WILSON, FRCS
A. AZMY, FRCS
Royal Hospital for Sick Children
Glasgow, UK.

1. Davies MS, Nadel S, Habibi P, Levin M, Hunt DM. The orthopaedic management of peripheral ischaemia in meningococcal septicaemia in children. *J Bone Joint Surg [Br]* 2000;82-B:383-6.
2. Farrar MJ, Bennet GC, Wilson NIL, Azmy A. The orthopaedic implications of peripheral limb ischaemia in infants and children. *J Bone Joint Surg [Br]* 1996;78-B:930-3.

Author's reply:

Sir,

We thank the correspondents for their interest in our article and their comments. We are certainly aware of the paper by Farrar et al which looked at peripheral gangrene from a number of causes

of which only one, in their series of 12 infants, was due to meningococcal septicaemia.

Our paper was not only focused on the management of ischaemia during meningococcal disease, but also, in particular, on the place of surgical treatment before the development of gangrene. Our main interest was to review the place of fasciotomy in the management of this condition, which was not discussed in their paper.

We agree that purpura fulminans is a consequence of thrombosis in small vessels, and that thrombectomy is not relevant. Certainly, heparin and tissue plasminogen activator have been used in these cases, but none of these agents affected the decision on whether or not to use a fasciotomy. Indeed, their use compounded the problems of fasciotomy, namely excessive bleeding and protein loss. We felt that the paper by Farrar et al looked at the effects of peripheral limb ischaemia rather than prevention, which was our aim.

D. HUNT, FRCS
St Mary's Hospital
London, UK.

Interstitial laser photocoagulation for the treatment of osteoid osteoma

Sir,

We read with interest the report in the November 2000 issue by Witt et al¹ entitled 'Interstitial laser photocoagulation for the treatment of osteoid osteoma: results of a prospective study'. We agree that open surgery for osteoid osteomas is no longer indicated. Their paper presents clear evidence of the effectiveness of percutaneous treatment.

We are concerned, however, that in laser photocoagulation the temperature cannot be accurately controlled. Also, the use of very fine-gauge needles makes biopsy impossible, and we note that none of the cases reported in this study had the diagnosis confirmed histologically.

In our unit, we have preferred to use CT-guided radiofrequency coagulation. This technique uses a trochar and cannula of a larger gauge (Bonopt; Bone Biopsy System, RADI Medical Systems), which allows material to be obtained for histological examination before the passage of the radiofrequency electrode into the lesion. This method also allows very precise regulation of temperature, which would have important advantages in anatomical sites close to other sensitive tissues, such as the spine.

P. COOL, FRCS Orth
D. H. WILLIAMS, MCh(Orth) FRCS
V. PULLICINO, FRCR
The Robert Jones and Agnes Hunt Orthopaedic & District Hospital NHS Trust, Shropshire, UK.

1. Witt JD, Hall-Craggs MA, Ripley P, Cobb JP, Brown SG. Interstitial laser photocoagulation for the treatment of osteoid osteoma: results of a prospective study. *J Bone Joint Surg [Br]* 2000;82-B:1125-8.

Authors' reply:

Sir,

We thank Mr Cool and his colleagues for their comments; they raise important issues. In patients with a typical history and radiological findings of osteoid osteoma we do not consider it necessary to obtain histological material. The cases presented in our paper were those in which the surgeon and bone-tumour radiologist were in agreement that the features were typical of

osteoid osteoma. Clearly, when there is doubt, a biopsy is indicated.

The issue of the temperatures reached in bone and the potential danger to surrounding normal structures is very important and in this there is little difference between interstitial laser photocoagulation (ILP) and radiofrequency-induced injury because of the nature of transmission of heat within bone.

The area of coagulative necrosis produced by ILP is controlled by the amount of energy delivered. Experimental work in bone has shown that the diameter of the necrosis around a fibre can be varied from 3.4 mm with 200 J to 9.2 mm with 1400 J.¹ The longitudinal area of coagulation is slightly more extensive. In the same study, thermocouples were inserted at varying distances from the fibre tip. The temperatures recorded at the tip were 90° to 280°C. At 10 mm from the tip, the temperature ranged from 37° to 56°C and did not increase significantly once the energy delivered had reached 1400 J.

These findings are similar to the experimental work of Tillotson, Rosenberg and Rosenthal,² investigating the thermal injury to bone produced by a radiofrequency probe with the probe temperature set to 80°C. The diameter of the necrosis ranged from 0.9 to 1.3 cm, and was found to be insensitive to varying durations of heat. No thermocouples were used in this study to document the temperature at varying distances from the probe, but the authors chose this temperature based on the work of Lundskog,³ who demonstrated the threshold for osteocyte necrosis to be 50°C for a duration of 30 seconds.

It is therefore true to say that the temperature at the tip of a radiofrequency probe can be precisely regulated. Since there is no practical means of measuring temperature in the bone beyond the probe in the clinical situation, however, the temperature and effect can only be inferred from experimental data.

Regardless of the technique used, great care is required when using a thermal modality close to neural elements and it is obviously important to be conversant with the specific physical properties of that particular technique.

J. D. WITT, FRCS Orth
M. A. HALL-CRAGGS, FRCR
The Middlesex Hospital and University College London Hospitals NHS Trust
London, UK.

1. Gangi A, Gasser B, De Unamuno S, et al. New trends in interstitial laser photocoagulation of bones. *Semin Musculoskelet Radiol* 1997;1:331-7.
2. Tillotson CL, Rosenberg AE, Rosenthal DI. Controlled thermal injury of bone: report of a percutaneous technique using radiofrequency electrode and generator. *Invest Radiol* 1989;24:888-92.
3. Lundskog J. Heat and bone tissue: an experimental investigation of the thermal properties of bone and threshold levels for thermal injury. *Scand J Plast Reconstr Surg* 1972;9:1-80.

Clinical outcome of congenital talipes equinovarus diagnosed antenatally by ultrasound

Sir,

I read with interest the article entitled 'Clinical outcome of congenital talipes equinovarus diagnosed antenatally by ultrasound'¹ in the August 2000 issue.

It is well known that congenital talipes equinovarus (CTEV) represents a spectrum, from so-called postural talipes requiring little active management, to a severe deformity requiring extensive surgery.² Assessment of severity, even after birth, remains difficult, with some authors classifying the condition on the basis of a response to a period of conservative management.³

Without accurate assessment of severity, the natural history of

the condition and the management required cannot be predicted. Indeed, 26% of those diagnosed antenatally by Tillett et al as having CTEV required no active management. These are presumably children with postural talipes as described by Porter, or even transient talipes as described by Bar-Hava et al.⁴ Given the high percentage of children requiring no treatment, it is difficult to justify the authors' statement that "when the appearances (on the antenatal scan) are abnormal, a significant abnormality is likely to be present". Other studies report a false-positive rate of up to 35%.⁵

The value of an antenatal diagnosis of talipes is in the fact that it may be associated with other genetic and morphological abnormalities and its presence therefore indicates the need for a detailed scan and karyotyping.⁶ Benacerraf⁷ found associated abnormalities in 15 of 18 fetuses with talipes diagnosed antenatally.

CTEV is a condition with a number of different causes which result in deformities of variable severity. For example, Macnicol and Nadeem⁸ have identified a subgroup of children with abnormalities of the somatosensory evoked potentials. Aetiological information of this type may ultimately prove to be of more value in predicting the natural history. Perhaps only when the aetiology of CTEV has been elucidated will it be possible to make a prediction of the natural history of the condition.

D. CHESNEY, FRCS Ed
Inverurie, Aberdeenshire, UK.

1. **Tillett RL, Fisk NM, Murphy M, Hunt DM.** Clinical outcome of congenital talipes equinovarus diagnosed antenatally by ultrasound. *J Bone Joint Surg [Br]* 2000;82-B:876-80.
2. **Porter RW.** Congenital talipes equinovarus: I. Resolving and resistant deformities. *J Bone Joint Surg [Br]* 1987;69-B:822-5.
3. **Hooker CH.** A clubfoot philosophy. *J Bone Joint Surg [Br]* 1985;67-B:490.
4. **Bar-Hava I, Bronshtein M, Orvieto R, et al.** Caution: prenatal clubfoot can be both a transient and a late-onset phenomenon. *Prenat Diagn* 1997;17:457-60.
5. **Woodrow N, Tran T, Umstad M, et al.** Mid-trimester ultrasound diagnosis of isolated talipes equinovarus: accuracy and outcome for infants. *Aust N Z J Obstet Gynaecol* 1998;38:301-5.

6. **Gonzalez P, Nicolini U, Rahman F, Fisk NM, Rodeck CH.** Does oligohydramnios cause fetal club foot? *J Obstet Gynaecol* 1991;11:11-4.
7. **Benacerraf BR.** Antenatal sonographic diagnosis of congenital club-foot: a possible indication for amniocentesis. *J Clin Ultrasound* 1986;14:703-6.
8. **Macnicol MF, Nadeem RD.** Evaluation of the deformity in club foot by somatosensory evoked potentials. *J Bone Joint Surg [Br]* 2000;82-B:731-5.

Authors' reply:

Sir;

We thank Mr Chesney for his comments on our paper. We agree that the severity of talipes is difficult to assess. Our point is that it is not possible to do this using antenatal ultrasound. Our statement that an abnormality detected on ultrasound correlates with talipes present at birth, was to stress that the purpose of this study was to help orthopaedic surgeons to counsel parents on whether or not their child has talipes and the likelihood of their baby needing treatment. We therefore stand by our statement, since these children all had talipes, although some cases were mild and needed only simple stretching. These were not normal feet, although they may well have been resolving postural talipes. It is important that surgeons are aware of these conditions when advising parents.

It is helpful to be able to tell parents that their child has a 26% chance of requiring only the simplest treatment and that the outcome will be excellent both functionally and cosmetically. The false-positive result described in our series was also important since it was part of a more complex deformity. We accept that other series have high false-positive rates but believe that these will reduce as antenatal scanning becomes more accurate.

Our paper was concerned only with the correlation between antenatal ultrasound and clinical outcome and not with the aetiology or natural history of talipes itself.

D. HUNT, FRCS
R. TILLET, MB, BS
St Mary's Hospital
London, UK.