Correspondence

We welcome letters to the Editor concerning articles which have recently been published. Such letters will be subject to the usual stages of selection and editing; where appropriate the authors of the original article will be offered the opportunity to reply.

Letters should normally be under 300 words in length, double-spaced throughout, signed by all authors and fully referenced. The edited version will be returned for approval before publication.

Reamed nailing of Gustilo grade-IIIB tibial fractures

Sir,

Successful management of Gustilo grade-IIIB fractures is an unsolved problem. I therefore read with interest the article in the November 2000 issue by Keating et al¹ entitled 'Reamed nailing of Gustilo grade-IIIB tibial fractures'.

There were, however, several inconsistencies in this article. Although 5% of their patients underwent 'early amputation' the authors did not describe the factors leading to amputation, except for mentioning the cause as "such was the severity of associated soft-tissue damage". Obviously, grade-IIIB fractures all have severe soft-tissue damage. It is very important to know which of these fractures required amputation. The algorithm presented in Figure 2 for the management of infection in Gustilo grade-IIIB fractures treated by reamed intramedullary nailing shows that the endpoint is invariably union, although amputation has to be undertaken in certain cases, and in their own study one patient underwent amputation. Albeit rare, this is too important an endpoint to be omitted. The authors mention that Keating et al² found that the only difference between reamed and unreamed nailing was the incidence of breakage of the implant. I checked the reference and found that this was actually breakage of screws. The authors should have clearly mentioned this, because breakage of the implant could also mean breakage of the nail. Most importantly, the authors mention that Schandelmaier et al³ and Tornetta et al⁴ who compared tibial nailing with external fixation for Gustilo grade-IIIB fractures, found that "with intramedullary nailing the time to union was shortened and the functional outcome improved". On checking the first reference, however, I found that Schandelmaier et al³ had actually concluded that the "Time to bony union, infection and nonunion were not significantly different between the groups". Moreover, on checking the article by Tornetta et al,⁴ I found that they had also clearly mentioned that "The differences in healing and range of motion were not statistically significant". Hence both of these references have been misrepresented. Finally, the authors conclude that "Our results indicate that intramedullary nailing is a satisfactory way of managing Gustilo grade-IIIB tibial fractures". In their study, however, 42% of patients had nonunion, 7% had malunion, 17.5% developed infection, 38% were unemployed with no likelihood of returning to work, 23.4% were too disabled for any recreational activity, 60% had residual ankle stiffness, 44.7% had persistent pain at the site of the fracture and 7% underwent amputation.

Based on these results it is not appropriate to conclude that reamed intramedullary nailing is a satisfactory treatment for Gustilo grade-IIIB tibial fractures. I believe that the conclusion should be that a prospective, randomised, case-control trial is required which compares reamed intramedullary nailing with other methods of management, such as the Ilizarov fixator.

V. SAHNI, MS, MCh Orth
Wirral Hospital NHS Trust
Wirral, UK.

1. Keating JF, Blachut PA, O'Brien PJ, Court-Brown CM. Reamed nailing of Gustilo grade-IIIB tibial fractures. *J Bone Joint Surg [Br]* 2000;82-B:1113-6.
2. Keating JF, O'Brien PJ, Blachut PA, Meek RN, Broekhuysen H. Locking intramedullary nailing with and without reaming for open fractures of the tibial shaft: a prospective, randomised study. *J Bone Joint Surg [Am]* 1997;79-A:334-41.
3. Schandelmaier P, Krettek C, Rudolf J, et al. Superior results of tibial rodding versus external fixation in grade 3B fractures. *Clin Orthop* 1997;342:164-72.
4. Tornetta P III, Bergman M, Watnik N, Berkowitz G, Steuer J. Treatment of grade IIIB open tibial fractures: a prospective, randomised comparison of external fixation and non-reamed locked nailing. *J Bone Joint Surg [Br]* 1994;76-B:13-9.

Author's reply:

Sir,

I thank Mr Sahni for his comments. The decision on whether to perform an early amputation in grade-IIIB open tibial fractures is a difficult one, and invariably comes down to a subjective assessment of the extent of soft-tissue loss and the likely functional outcome. The decision to amputate in three cases at an early stage was made by experienced orthopaedic trauma surgeons, usually in conjunction with plastic surgeons. We did not routinely use any of the scoring systems for limb damage to assist in the decision since none of the proposed scores has been shown to be reliable.

The algorithm which we proposed for the management of infection did not include amputation because the aim of treating infection after these injuries is to eradicate the infection. Amputation represents failure to achieve this aim and the algorithm was designed with the avoidance of this endpoint in mind.

We consider the literature to support the view that functional outcome is better with intramedullary nailing than with external fixation for these injuries. The results which we recorded in terms of complications and functional outcome are very similar to other reports in the literature. The need for further surgical intervention is common in these injuries irrespective of which method of treatment is used, and after consideration we see no reason to alter our conclusion that reamed nailing is satisfactory for these fractures. We agree with Mr Sahni that a prospective trial comparing reamed nailing with another method of treatment would be a useful addition to the literature, although we would not agree that the Ilizarov fixator would be the obvious alternative. There are no published results on the use of this device in the acute management of these injuries, and the complex soft-tissue problems associated with grade-IIIB open fractures make the use of any external fixator difficult.

J. KEATING, MPhil, FRCS Ed(Orth)
Edinburgh Royal Infirmary
Edinburgh, UK.

Radiation therapy for the prevention of heterotopic ossification at the elbow

Sir,

We read with interest the article in the April 2001 issue entitled 'Radiation therapy for the prevention of heterotopic ossification (HO) at the elbow' by Heyd et al¹. We agree with the authors that prophylactic radiotherapy can be effective in reducing the recurrence of HO after surgery. We were, however, surprised to find that the authors failed to mention the timing of the surgery from the onset of the formation of heterotopic bone.

Garland et al² have recommended delaying surgery for 12 to 18 months from the onset of the formation of heterotopic bone to reduce the incidence of recurrence. This is effective, but at the expense of the range of movement of the elbow resulting in poor final function. Early surgical excision of heterotopic bone has the advantage of preserving the range of movement and, when combined with radiotherapy, can significantly reduce the rate of recurrence.³

In view of these findings, we believe that the timing of surgery has a strong influence on the rate of recurrence after excision of HO. Without this information it would be inappropriate to attribute the absence of any recurrence entirely to the effect of radiotherapy.

S. ANKARATH, FRCS Ed, FRCS (Tr & Orth)
P. V. GIANNOUDIS, MD
St James's University Hospital
Leeds, UK.

1. Heyd R, Strassmann G, Schopohl B, Zamboglou N. Radiation therapy for the prevention of heterotopic ossification at the elbow. *J Bone Joint Surg [Br]* 2001;83-B:332-4.
2. Garland DE, Hanscom DA, Keenan MA, Smith C, Moore T. Resection of heterotopic ossification in the adult with head trauma. *J Bone Joint Surg [Am]* 1985;67-A:1261-9.
3. McAuliffe JA, Wolfson AH. Early excision of heterotopic ossification about the elbow followed by radiation therapy. *J Bone Joint Surg [Am]* 1997;79-A:749-55.

Authors' reply:

Sir,

Mr Ankarath and Dr Giannoudis mentioned in their letter that the delay between the radiological manifestation of heterotopic ossification (HO) and the time of the surgical excision has been reported to be a prognostic factor for local recurrence in neurogenic HO after head trauma.¹ In our group of patients, only one patient had neurogenic HO, one had spontaneous development of HO and the remaining seven had traumatic HO. We agree with Garland² that the type of HO is relevant when considering the time of surgical excision. Traumatic HO may be resected six months after the injury, whereas for neurogenic HO (traumatic brain injury or spinal cord injury) 12 to 18 months is necessary.²

In our group of patients these recommendations were followed. In the seven patients with traumatic HO and in the one with spontaneous HO, excision was performed six to eight months after diagnosis, whereas in the patient with neurogenic HO (cerebral bleeding), it was carried out 12 months after the injury when the patient had sufficient relief of spasticity.

From our clinical experience, HO at the elbow commonly causes severe limitation of movement of the joint, and in order to shorten the period of discomfort for the patient, excision for traumatic HO should be performed after six months. In cases of

neurogenic HO, a longer delay is needed and the decision on the timing of surgery depends on the clinical findings with regard to the relief of neurological deficits.

R. HEYD, MD
Klinikum Offenbach
Offenbach, Germany.
G. STRASSMANN, MD
University Hospital
Marburg, Germany.

1. Garland DE, Hanscom DA, Keenan MA, Smith C, Moore T. Resection of heterotopic ossification in the adult with head trauma. *J Bone Joint Surg [Am]* 1985;67-A:1261-9.
2. Garland DE. A clinical perspective on common forms of acquired heterotopic ossification. *Clin Orthop* 1991;263:13-29.

Conservative versus operative treatment for displaced ankle fractures in patients over 55 years of age

Sir,

I read with interest the article in the May 2001 issue entitled 'Conservative versus operative treatment for displaced ankle fractures in patients over 55 years of age.'¹ I thank the authors for establishing the fact that operative treatment is better than conservative treatment for such fractures. I wish, however, to bring to their attention a few points for discussion.

The type of fracture (Weber or AO types and subtypes) is not mentioned which makes it difficult to interpret the results. Weber C fractures, although less common in this age group, are difficult to treat conservatively, which could influence the final result.

Most of the displaced fractures of the ankle involve a rotational component (PER and SER) and require an initial above-knee cast for at least four weeks.² The authors have used a below-knee cast only for conservative management.

Although Fernandez³ and Ali et al⁴ have preferred operative to conservative treatment in this age group, in the form of posterolateral plating or a dynamic compression plate, respectively, they also reported recognised wound complications of 6%⁴ and removal of the implant in 14%. Winkler, Weber and Simpson⁵ preferred posterolateral plating to conservative treatment, but found poor results in 10% of elderly patients when compared with the young. The authors state that in the operative group two patients had wound problems, two had a complex regional pain syndrome, two had tenderness of the scar and in four the implants were removed, although they had better overall results than the conservative management group as measured using the Olerud score. It should be noted that the Olerud Molander scoring system has no negative points for the above complications, and no significant difference has been reported among the visual analogue scores in both groups. This study has included various forms of internal fixation such as plating, Rush nailing and K-wire fixation which could have influenced the result. The article does not specify the choice of implant, which is crucial in this age group.

P. R. RAMASAMY, FRCS, MS Orth
Weston General Hospital
Weston Super Mare, UK.

1. Makwana NK, Bhowal B, Harper WM, Hui AW. Conservative versus operative treatment for displaced ankle fractures in patients over 55 years of age. *J Bone Joint Surg [Br]* 2001;83-B:525-9.
2. Rockwood CA, et al. Fractures and injuries of the ankle. In: Rockwood CA, Green DP, Bucholz RW. *Fractures in adults*. 4th edition, Vol. 2. Philadelphia: J. B. Lippincott, 1996:2226.
3. Fernandez GN. Internal fixation of the oblique, osteoporotic fracture of the lateral malleolus. *Injury* 1988;19:257-8.

4. Ali MS, McLaren CAN, Rouholamin E, O'Connor BT. Ankle fractures in the elderly: nonoperative or operative treatment. *J Orthop Trauma* 1987;1:275-80.
5. Winkler B, Weber BG, Simpson LA. The dorsal antiglide plate in the treatment of Danis-Weber type-B fractures of the distal fibula. *Clin Orthop* 1990;259:204-9.

Authors' reply:

Sir,

We thank Mr Ramasamy for his interest in our paper. We classified our fractures using the AO system. There were 18 Weber B and four Weber C fractures in the open reduction and internal fixation (ORIF) group, and 14 Weber B and seven Weber C fractures in the closed treatment group. There was no statistically significant difference ($p=0.263$) between the two groups. While Weber C fractures usually involve damage to the syndesmotic ligament (similar to a pronation/eversion injury), Weber B fractures may also involve the anterior and posterior tibiofibular ligaments, as well as the syndesmotic ligament (supination/external rotation or pronation/abduction). These are just as difficult to treat conservatively as Weber C fractures, and we would disagree that the distribution of fractures influenced our results. Our trial was planned to take into consideration the reducibility of the fractures and all patients started at the same point, i.e., the fracture was reduced according to our protocol before being randomised into the study.

While there may be a theoretical advantage for treating rotational injuries of the ankle with an above-knee plaster cast, we have not seen any evidence or prospective randomised studies to support this. Our study was based on the standard practice in Leicester of the management of fractures of the ankle and this probably represents that followed in many other units. An above-knee plaster cast in an elderly patient is likely to impair mobility significantly and may also increase the risk of complications such as stiffness of the knee. In addition, some elderly patients cannot cope with this type of cast.

Overall, there was a higher rate of complications observed in the closed treatment group than in the ORIF group if loss of reduction is included. Most of the complications in the ORIF group resolved by three months, with one patient awaiting removal of the implant and two who had tenderness of the scar at review. While no scoring system is ideal, the composite result using the Olerud score, visual analogue score, range of movement of the ankle and ankle girth probably reflects the outcome better than any one individual score.

The aim of open reduction and internal fixation in all cases was to achieve anatomical reduction and stable fixation. Nearly all the fractures of the ankle (96%) were fixed using a 1/3 semitubular plate on the lateral side, with or without lag screws. In one patient the lateral malleolus was treated by a Rush pin with a good result. AO cancellous screws were used in 82% of cases on the medial side, and this was supplemented in one by a Kirschner wire. No fixation was necessary in three medial malleoli.

N. K. MAKWANA, FRCS Ed (Tr & Orth)
B. BHOWAL, FRCS (Tr & Orth)
W. M. HARPER, MD
University Hospitals of Leicester NHS Trust
Leicester, UK.
A. W. HUI, FRCS (Tr & Orth)
Middlesborough General Hospital
Middlesborough, UK.

et al¹ entitled 'Conservative *versus* operative treatment for displaced ankle fractures in patients over 55 years of age'. Within the study, all conservatively treated fractures of the ankle were managed by the application of a moulded below-knee plaster cast.

It is well known that in order to control rotation at the ankle, a cast needs to be extended to above the knee with the knee in slight flexion.^{2,3} Fractures associated with syndesmotic disruption (pronation/external rotation, abduction/external rotation, or Weber type C) are rotationally unstable, and if they are to be managed conservatively, an above-knee cast is indicated.

The authors defined the fractures within the study using the AO classification system, and by the number of malleoli involved, and stated that in the conservatively managed group, there was no significant difference in the type of fracture between the patients in whom there was some loss of reduction and those in whom there was none. The Lauge-Hansen classification, however, gives a clearer definition based on the aetiology of the fracture, and with this system it is easier to predict potential rotational instability. Given the relatively small numbers of patients within each of the groups in the study, and the classification systems used, the significance of an increased loss of reduction in rotationally unstable fractures may well not have been apparent.

Below-knee casts were used for all conservatively treated fractures of the ankle which, as stated above, is not appropriate management for rotationally unstable fractures. These fractures may well appear to be well reduced initially, but if treated with a below-knee cast, will have a propensity to displace after the initial radiographs. The failure to address this aspect of the conservative treatment of fractures of the ankle may have influenced the outcome in the closed treatment group, in what was otherwise a very well constructed trial.

I. D. McDERMOTT, FRCS
Imperial College
London, UK.

1. Makwana NK, Bhowal B, Harper WM, Hui AW. Conservative *versus* operative treatment for displaced ankle fractures in patients over 55 years of age. *J Bone Joint Surg [Br]* 2001;83-B:525-9.
2. Geissler WB, Tsao AK, Hughes JL. Fractures and injuries of the ankle. In: Rockwood CA Jr, Green DP, Bucholz RW, Heckman JD, eds. *Fractures in adults*. Philadelphia: Lippincott-Raven, 1996:2226.
3. Sean-Hughes PF. External rotational injury of the ankle joint with displacement of the talus. *Ann R Coll Surg Engl* 1977;59:61-4.

Authors' reply:

Sir,

We thank Mr McDermott for his interest in our paper. We classified our fractures according to the AO system. Although the Lauge-Hansen classification system characterises the mechanism and sequence of injury, it is more complex and not all fractures conform exactly to one of the patterns described.¹ It is difficult to apply in a reproducible way,² and the inter- and intraobserver agreement is poor with the more common types of fracture.³

With regard to the use of above-knee plaster casts in elderly patients we would draw your attention to our replies to Mr Fowler and Mr Ramasamy.

N. K. MAKWANA, FRCS Ed (Tr & Orth)
B. BHOWAL, FRCS (Tr & Orth)
W. M. HARPER, MD
University Hospitals of Leicester NHS Trust
Leicester, UK.
A. W. HUI, FRCS (Tr & Orth)
Middlesborough General Hospital
Middlesborough, UK.

1. Yde J. The Lauge-Hansen classification of malleolar fractures. *Acta Orthop Scand* 1980;51:181-92.

Sir,

I read with interest the article in the May 2001 issue by Makwana

2. **Nielsen JO, Dons-Jensen H, Sorensen HT.** Lauge-Hansen classification of malleolar fractures: an assessment of the reproducibility in 118 cases. *Acta Orthop Scand* 1990;61:385-7.
3. **Thomsen NO, Overgaard S, Olsen LH, Hansen H, Nielsen ST.** Observer variation in the radiographic classification of ankle fractures. *J Bone Joint Surg [Br]* 1991;73-B:676-8.

Sir,

I read with interest the article by Makwana et al¹ in the May 2001 issue entitled 'Conservative *versus* operative treatment for displaced ankle fractures in patients over 55 years of age'.

This Leicester-controlled trial revealed a loss of reduction of 38% in patients in the conservative group who were treated by below-knee casts. This is not surprising, since most displaced fractures of the ankle are caused by external rotation violence. It is very difficult to control rotation with a below-knee cast because the upper end of the plaster loosens as the swelling decreases, allowing the foot to rotate externally (the Wellington-boot phenomenon).

One way to control rotation is to apply an above-knee plaster with the knee at 90° of flexion, but this plaster is very cumbersome and prevents weight-bearing. A better solution is to use an above-knee plaster combined with the Bridgend knee grip which contours the plaster closely around the knee. The technique is simple. After reduction and the application of a below-knee cast, the foot is strongly internally rotated by an assistant and the plaster is extended to the mid-thigh. It is then closely moulded around the sides of the patella using the radial sides of both thumbs, and around the femoral condyles using the fingers (Fig. 1).

Unlike internal fixation, this method of controlling deformity due to rotation allows full weight-bearing in the plaster at an early stage. The lower limb is designed for weight-bearing, which assists in the nutrition of bone and cartilage and prevents subsequent stiffness of the ankle and sympathetic osteodystrophy.

There are many areas in the world where operative fixation of fractures is not feasible, and we therefore need to continue to promote adequate methods of conservative treatment.

A. W. FOWLER, FRCS
Bridgend, Mid Glamorgan, UK.

1. **Makwana NK, Bhowal B, Harper WM, Hui AW.** Conservative *versus* operative treatment for displaced ankle fractures in patients over 55 years of age. *J Bone Joint Surg [Br]* 2001;83-B:525-9.

Fig. 1

Diagram showing the application of the above-knee plaster and Bridgend knee grip.

2. **Fowler AW.** The importance of joint movements in the treatment of fractures. *East Central Afr J Surg* 1995;1:49-60.

Authors' reply:

Sir,

We thank Mr Fowler for his comments regarding the treatment of rotationally unstable fractures of the ankle with an above-knee plaster cast combined with the Bridgend knee grip. We have had no experience with the grip and cannot comment on its use. While we agree there is a theoretical advantage of using an above-knee cast, we have not seen any evidence or prospective, randomised studies to support its use over a below-knee cast. It may be justified in areas where access to surgical facilities is limited; the plaster suggested would form a useful addition to the armamentarium when open reduction and internal fixation is not a feasible option. In developing countries where access is not an issue, however, we should aim to restore anatomical congruity of the ankle in the best way possible to achieve the best possible results. In this respect our study supports open reduction and internal fixation of these fractures.

N. K. MAKWANA, FRCS Ed (Tr & Orth)
B. BHOWAL, FRCS (Tr & Orth)
W. M. HARPER, MD
University Hospitals of Leicester NHS Trust
Leicester, UK.
A. W. HUI, FRCS (Tr & Orth)
Middlesbrough General Hospital
Middlesbrough, UK